

Stacey (SW 11-2363).

MARKET SEGMENT

R

Retail
Baby potatoes

Early maincrop red skin small tuber size ware variety with a high tuber set and a yellow flesh colour.

- ▶ Oval round small sized
- ▶ Firm cooking
- ▶ Good tolerance against bruising and damage



CHARACTERISTICS

General characteristics

Breeder: Lantmännens Seed B.V.
Crossing: ROSAGOLD x AR 04-3064
Seedling: SW 11-2363
Breeder rights: Pending

Plant and tuber characteristics

Maturity: maincrop / 5
Light sprout colour: -
Flower colour: light purple
Number of berries: none / 4
Foliage: - initial development: quite fast / 7,5
Foliage: - final development: good / 8
Foliage: - selectability: average / 6,5
Foliage: - plant firmness: quite firm / 7

Flesh colour: light yellow / 6,5
Skin colour: red
Skin finish: good / 7
Tuber size: average / 6,5
Tuber shape: oval round
Regularity of tuber shape: regular / 7
Tuberisation: high / 8,5
Regularity size: quite regular / 6,5

Sensitivity and quality

Dormancy: average / 6
Metribuzin: - / -
Bentazon: - / -
Ethylene: - / -

Secondary growth: quite sensitive / 6,5
Growth cracks: slightly sensitive / 7
Black spot (internal bruising): slightly sensitive / 7
Harvest damage: slightly sensitive / 7,5

Underwater weight: 423
Dry matter content: 22,8%
Dry matter distribution: good / 7,5
TGA - level: -

Crisping quality: unsuitable / 4
Fries quality: unsuitable / 4
Cooking type: rather firm / AB
Raw discolouration: quite sensitive / 6
Discolouration after cooking: slightly sensitive / 7

HOW TO GROW

Fertilization¹

Nitrogen: standard advice.

Phosphate: standard advice.

Potassium: standard advice.

¹ Level of fertilization is based on soil analysis

Seed treatment

Desprouting more than once can have a negative impact on yield and size regularity.

Planting distance

28/35 mm: 22 cm (60,500 plants/ha)

35/55 mm: 27 cm (50,000 plants/ha)

Weeds

Metribuzin (Sencor) scores are not yet determined.

Fungicides

A preventive spraying schedule against late blight is advised.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

Storage

Preferred storage temperature is 5 °C. Suitable for medium term storage.



Say potato,
say **Agrico.**

Resistances

Golden nematode Ro 1/4: resistant / 9

Golden nematode Ro 2/3: - / -

White nematode Pa 2: - / -

White nematode Pa 3: - / -

Late blight-foilage: slightly susceptible / 7

Late blight-tuber: slightly susceptible / 6

Common scab: susceptible / 5

Powdery scab: - / -

Fusarium: very susceptible / 4

Erwinia: - / -

Black dot: - / -

Spraing: slightly susceptible / 6

Virus - Leaf Roll: - / -

Virus - A: - / -

Virus - X: - / -

Virus - Yn: very susceptible / 4

Virus - Yntn: - / -

Wart disease F1: resistant* / 10*

Wart disease F6: - / -

Wart disease F18: - / -

* Results marked with an asterisk (*) are based on our own research, and are not yet officially published