


Rousseau.

MARKET SEGMENT



A second early maincrop red skin salad variety with a beautiful shape and a very regular size.

- ▶ High tuberisation
- ▶ Nice skin finish
- ▶ Official registration for chair ferme



CHARACTERISTICS

General characteristics

Breeder: Lantmännen Seed B.V.
Crossing: CECILE x ROSAGOLD
Seedling: SW 08-0012
Breeder rights: 31-12-2049

Plant and tuber characteristics

Maturity: second early / 6,5
Light sprout colour: red violet
Flower colour: purple
Number of berries: average / 6
Foliage: - initial development: fast / 8
Foliage: - final development: good / 8
Foliage: - selectability: quite easy / 7
Foliage: - plant firmness: quite firm / 7

Flesh colour: light yellow / 6,5

Skin colour: red

Skin finish: good / 7

Tuber size: quite large / 7

Tuber shape: long oval

Regularity of tuber shape: regular / 7

Tuberisation: high / 8

Regularity size: quite regular / 6,5

Sensitivity and quality

Dormancy: average / 6

Metribuzin: quite sensitive / 6

Bentazon: - / -

Ethylene: 5% fewer tubers / 4

Secondary growth: sensitive / 5,5

Growth cracks: slightly sensitive / 7

Black spot (internal bruising): slightly sensitive / 7,5

Harvest damage: slightly sensitive / 7,5

Underwater weight: 378

Dry matter content: 20,6%

Dry matter distribution: - / -

TGA - level: 5,9

Crisping quality: unsuitable / 4

Fries quality: unsuitable / 4,5

Cooking type: rather firm / AB

Raw discolouration: sensitive / 5

Discolouration after cooking: slightly sensitive / 7



www.agrico.nl/en/rousseau

HOW TO GROW

Fertilization¹

Nitrogen: standard advice.

Phosphate: standard advice.

Potassium: standard advice + 20%.

¹ Level of fertilization is based on soil analysis

Seed treatment

Desprouting can have a negative impact on yield and size regularity. Can be planted directly out of the cold store, preferably after a short heat shock.

Planting distance

30/40 mm: 20-22 cm (65,000 plants/ha)

40/50 mm: 22-26 cm (55,000 plants/ha)

Weeds

After emerge, a standard use of metribuzin (Sencor) can give some foliage reaction, but rarely any yield reduction.

Fungicides


A strict preventive spraying schedule against late blight is necessary.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

Storage

Preferred storage temperature is 4 °C. Suitable for long term storage.



Say potato,
say **Agrico.**

Resistances

Golden nematode Ro 1/4: resistant / 9

Golden nematode Ro 2/3: - / -

White nematode Pa 2: - / -

White nematode Pa 3: - / -

Late blight-foliage: very susceptible / 4,5

Late blight-tuber: susceptible / 5

Common scab: slightly susceptible / 6

Powdery scab: slightly susceptible / 6,5

Fusarium: slightly susceptible / 6,5

Erwinia: - / -

Black dot: slightly to not susceptible / 8

Spraing: very susceptible / 4,5

Virus - Leaf Roll: - / -

Virus - A: - / -

Virus - X: - / -

Virus - Yn: slightly susceptible / 7,5

Virus - Yntn: slightly susceptible / 6

Wart disease F1: - / -

Wart disease F6: - / -

Wart disease F18: - / -