

Jamaica (AR 14-0334).

An early traditional and retail variety with a good skin finish and size.




Retail
Fresh



Traditional

- ▶ **Good yield**
- ▶ **Good skin finish**
- ▶ **Good size**



characteristics

General characteristics

Breeder: Agrico Research B.V.

Crossing: OS 07-0020 x SAVIOLA

Breedersright: Pending

Seedling: AR 14-0334

Plant and tuber characteristics

Maturity: early / 7,5

Light sprout colour: - / -

Flower colour: white / W

Number of berries: above average / 5

Foliage - initial development: fast / 8

Foliage - final development: quite good / 7,5

Foliage - firmness: average / 6,5

Skin colour: yellow / Y

Flesh colour: light yellow / 6,5

Skin finish: good / 7

Tuber shape: oval long / OL

Regularity of tuber shape: regular / 7

Tuberisation: average / 6

Tuber size: large / 8

Regularity of size: regular / 7

Dormancy: quite short / 5,5

Sensitivity and quality

Secondary growth: quite sensitive / 6,5

Black spot (internal bruising): slightly to not sensitive / 8

Growth cracks: quite sensitive / 6,5

Harvest damage: quite sensitive / 6

Metribuzin - pre-emergence: very sensitive / 3

Metribuzin - post-emergence: very sensitive / 3

Ethylene: - / -

Underwater weight: 332

Dry matter content: 18,3%

Cooking type: rather firm / AB

Discolouration (after cooking): - / -

Raw discolouration: - / -

Fries quality: - / -

Dry matter distribution: - / -

Crisping quality: - / -

TGA-level: 15

How to grow



www.agricopotatoes.com

Fertilization ¹

Nitrogen: standard advice. Preferably as split application, second treatment shortly after tuberization.

Phosphate: standard advice.

Potassium: standard advice.

Seed treatment

A short heat shock or proper pre-sprouting is recommended.

Planting distance

28/35 mm: 20 cm (67.000 plants/ha)

35/50 mm: 27 cm (50.000 plants/ha)

Weeds

Do not use metribuzin (Sencor), as it gives a severe foliage reaction and a serious yield reduction.

Fungicides

A strict preventive spraying schedule against late blight is necessary.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

Storage - .

Preferred storage temperature is 4 °C. Suitable for delivery straight from the field or short term storage.

¹ Level of fertilization is based on soil analysis.



Say potato,
say **Agrico.**

Resistances

Virus - Leaf roll: - / -

Virus - Yn: slightly to not susceptible / 8

Virus - Yntn (tuber): slightly susceptible / 6

Spraing: slightly susceptible / 6,5

Late blight - foliage: very susceptible / 3,5

Late blight - tuber: slightly susceptible / 6

Fusarium: - / -

Common scab: susceptible / 5,5

Powdery scab: - / -

Black dot: - / -

Wart disease F1: resistant* / 10*

Wart disease F6: - / -

Wart disease F18: - / -

Golden nematode Ro 1/4: susceptible* / 1*

Golden nematode Ro 2/3: - / -

White nematode Pa 2: - / -

White nematode Pa 3: - / -