CULTIVATION RECOMMENDATION


Jamaica (AR 14-0334).

An early traditional and retail potato variety with a good skin finish and size.

Retail
Fresh

Traditional

- **Good yield**
- **Good skin finish**
- **Good size**



characteristics

General characteristics

Breeder: Agrico Research B.V.
Crossing: OS 07-0020 x SAVIOLA
Breedersright: Pending
Seedling: AR 14-0334

Plant and tuber characteristics

Maturity: early / 7,5
Light sprout colour: - / -
Flower colour: white / W
Number of berries: above average / 5
Foliage - initial development: fast / 8
Foliage - final development: quite good / 7,5
Foliage - firmness: average / 6,5
Skin colour: yellow / Y
Flesh colour: light yellow / 6,5
Skin finish: good / 7
Tuber shape: oval long / OL
Regularity of tuber shape: regular / 7
Tuberisation: average / 6
Tuber size: large / 8
Regularity of size: regular / 7
Dormancy: quite short / 5,5

Sensitivity and quality

Secondary growth: quite sensitive / 6,5
Black spot (internal bruising): slightly to not sensitive / 8
Growth cracks: quite sensitive / 6,5
Harvest damage: quite sensitive / 6
Metribuzin - pre-emergence: very sensitive / 3
Metribuzin - post-emergence: very sensitive / 3
Ethylene: - / -
Underwater weight: 332
Dry matter content: 18,3%
Cooking type: rather firm / AB
Discolouration (after cooking): quite sensitive / 6
Raw discolouration: - / -
Fries quality: - / -
Dry matter distribution: - / -
Crisping quality: - / -
TGA-level: -

Agrico has taken all reasonable care in producing this variety description. However, please note that any information may be subject to change and Agrico accepts no responsibility for errors or omissions. In no event shall Agrico be liable for damages arising out of or related to the information contained within this description.

AGRICO

CULTIVATION RECOMMENDATION

How to grow

www.agricopotatoes.com

Fertilization [1]

Nitrogen: standard advice.Preferably as split application, second treatment shortly after tuberization.

Phosphate: standard advice.

Potassium: standard advice.

Seed treatment

A short heat shock or proper pre-sprouting is recommended.

Planting distance

28/35 mm: 23 cm (58.000 plants/ha)

35/55 mm: 30 cm (44.500 plants/ha)

Weeds

Do not use metribuzin (Sencor), as it gives a severe foliage reaction and a serious yield reduction.

Fungicides

A strict preventive spraying schedule against late blight is necessary.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

Storage - .

Preferred storage temperature is 4 ºC. Suitable for delivery straight from the field or short term storage.

[1] Level of fertilization is based on soil analysis.



Say potato, say Agrico.

Resistances

Virus - Leaf roll: - / -

Virus - Yn: slightly to not susceptible / 8

Virus - Yntn (tuber): slightly susceptible / 6

Spraing: slightly susceptible / 6,5

Late blight - foliage: very susceptible / 3,5

Late blight - tuber: slightly susceptible / 6

Fusarium: - / -

Common scab: susceptible / 5,5

Powdery scab: - / -

Black dot: - / -

Wart disease F1: resistant / 10

Wart disease F6: - / -

Wart disease F18: - / -

Golden nematode Ro 1/4: susceptible* / 1*

Golden nematode Ro 2/3: - / -

White nematode Pa 2: - / -

White nematode Pa 3: - / -

Agrico • Duit 15 • 8305 BB Emmeloord • The Netherlands +31 (0)527 63 99 11 info@agrico.nl www.agricopotatoes.com