

Feliz.

A second early, salad variety with oval long tubers, cooking type A and strong resistance against PVY.



**Retail
Salads**

Image:

- ▶ **Good skin finish**
- ▶ **Cooking type A**
- ▶ **Good PVY resistance**

characteristics

General characteristics

Breeder: Lantmännen Seed B.V.
 Crossing: LORELEY x SW 06-0296
 Breedersright: Pending
 Seedling: SW 14-7008

Plant and tuber characteristics

Maturity: second early / 7
 Light sprout colour: - / -
 Flower colour: white / W
 Number of berries: - / -
 Foliage - initial development: fast / 8
 Foliage - final development: good / 8
 Foliage - firmness: quite firm / 7
 Skin colour: yellow / Y
 Flesh colour: light yellow / 6,5
 Skin finish: good / 7
 Tuber shape: oval long / OL
 Regularity of tuber shape: quite regular / 6,5
 Tuberisation: high / 8
 Tuber size: average / 6,5
 Regularity of size: quite regular / 6,5
 Dormancy: average / 6

Sensitivity and quality

Secondary growth: quite sensitive / 6
 Black spot (internal bruising): slightly to not sensitive / 8
 Growth cracks: slightly sensitive / 7
 Harvest damage: slightly to not sensitive / 8
 Metribuzin - pre-emergence: slightly sensitive / 7,5
 Metribuzin - post-emergence: slightly sensitive / 7,5
 Ethylene: 5% to 15% more tubers / 6
 Underwater weight: 358
 Dry matter content: 19,6%
 Cooking type: firm / A
 Discolouration (after cooking): - / -
 Raw discolouration: very sensitive / 4
 Fries quality: - / -
 Dry matter distribution: - / -
 Crisping quality: - / -
 TGA-level: -

How to grow



www.agricopotatoes.com

Fertilization ¹

Nitrogen: standard advice. Preferably in three applications, with second treatment shortly after tuberization.

Phosphate: standard advice.

Potassium: standard advice.

Seed treatment

A short heat shock or proper pre-sprouting is recommended. Desprouting can have a negative impact on yield and size regularity.

Planting distance

30/50 mm: 20 cm (67.000 plants/ha)

Weeds

A standard use of metribuzin (Sencor), before or after emerge, can give some foliage reaction, but rarely any yield reduction.

Fungicides

A strict preventive spraying schedule against late blight is necessary.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

Storage -

Preferred storage temperature is 4 °C. Suitable for medium term storage.

¹ Level of fertilization is based on soil analysis.

Image:

Say potato,
say Agrico.

Resistances

Virus - Leaf roll: - / -

Virus - Yn: slightly susceptible / 7

Virus - Yntn (tuber): - / -

Spraing: very susceptible / 4

Late blight - foliage: very susceptible / 3,5

Late blight - tuber: susceptible / 5,5

Fusarium: - / -

Common scab: susceptible / 5

Powdery scab: - / -

Black dot: - / -

Wart disease F1: resistant* / 10*

Wart disease F6: - / -

Wart disease F18: - / -

Golden nematode Ro 1/4: resistant / 9

Golden nematode Ro 2/3: susceptible* / 4*

White nematode Pa 2: moderately susceptible* / 5*

White nematode Pa 3: susceptible* / 3*