CULTIVATION RECOMMENDATION

Carolus.

An early maincrop, multipurpose ware potato variety with attractive red eyes and resilient to late blight in foliage and tuber.

- **Multipurpose variety**
- **High dry matter content**
- **Floury cooking**

Organic **Retail** Fresh **Traditional**



characteristics

General characteristics

Breeder: Agrico Research B.V.
Crossing: AGRIA x AR 00-94-17
Breedersright: 31-12-2043
Seedling: AR 02-3225

Plant and tuber characteristics

Maturity: early maincrop / 6
Light sprout colour: blue violet / BV
Flower colour: purple / P
Number of berries: very high / 4
Foliage - initial development: average / 6
Foliage - final development: good / 8
Foliage - firmness: firm / 8
Skin colour: yellow with red eyes / Y/R
Flesh colour: yellow / 7
Skin finish: good / 7
Tuber shape: oval / O
Regularity of tuber shape: regular / 7
Tuberisation: average / 6
Tuber size: quite large / 7
Regularity of size: regular / 7
Dormancy: average / 6,5

Sensitivity and quality

Secondary growth: slightly sensitive / 7
Black spot (internal bruising): quite sensitive / 6
Growth cracks: sensitive / 5,5
Harvest damage: sensitive / 5,5
Metribuzin - pre-emergence: very sensitive / 4,5
Metribuzin - post-emergence: quite sensitive / 6
Ethylene: 15% to 25% more tubers / 7
Underwater weight: 398
Dry matter content: 21,6%
Cooking type: floury / BC
Discolouration (after cooking): quite sensitive / 6
Raw discolouration: slightly sensitive / 7
Fries quality: good / 7
Dry matter distribution: excellent / 8
Crisping quality: good / 7
TGA-level: 5,4 mg/100g

Agrico has taken all reasonable care in producing this variety description. However, please note that any information may be subject to change and Agrico accepts no responsibility for errors or omissions. In no event shall Agrico be liable for damages arising out of or related to the information contained within this description.

AGRICO

CULTIVATION RECOMMENDATION

How to grow

www.agricopotatoes.com

Fertilization [1]

Nitrogen: standard advice + 10%.
Phosphate: standard advice.
Potassium: standard advice + 15%.

Seed treatment

A short heat shock or proper pre-sprouting is recommended.Preferably plant the tubers into warm soil, as the initial development of this variety is slow.Desprouting can have a negative impact on yield and size regularity.

Planting distance

28/35 mm: 20 cm (66.500 plants/ha)
35/55 mm: 27 cm (49.000 plants/ha)

Weeds

Do not use metribuzin (Sencor), as it gives a severe foliage reaction and a serious yield reduction.

Fungicides

Despite the late blight resistance, remain alert for infections.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

Storage - .

Preferred storage temperature is 5 ºC. For processing purposes, do not store below 7 ºC. Suitable for medium term storage.

[1] Level of fertilization is based on soil analysis.



Say potato, say Agrico.

Resistances

Virus - Leaf roll: - / -
Virus - Yn: slightly susceptible / 7
Virus - Yntn (tuber): slightly susceptible / 7
Spraing: slightly susceptible / 6
Late blight - foliage: slightly to not susceptible / 9
Late blight - tuber: slightly to not susceptible / 9
Fusarium: slightly susceptible / 6
Common scab: susceptible / 5
Powdery scab: susceptible / 5
Black dot: slightly susceptible / 6
Wart disease F1: resistant / 10
Wart disease F6: - / -
Wart disease F18: - / -
Golden nematode Ro 1/4: - / -
Golden nematode Ro 2/3: - / -
White nematode Pa 2: - / -
White nematode Pa 3: - / -

Agrico • Duit 15 • 8305 BB Emmeloord • The Netherlands +31 (0)527 63 99 11 info@agrico.nl www.agricopotatoes.com