

Cerisa.

An early, small tuber sized red skin ware and salad speciality potato with an excellent flavour.



**Retail
Salads**

- ▶ **Very high tuberisation**
- ▶ **Excellent flavour and cooking properties**
- ▶ **Good virus resistances**



characteristics

General characteristics

Breeder: Desmazières

Crossing: FRANCINE x LAURA

Breedersright: 31-12-2041

Seedling: DM 99-063-07

Plant and tuber characteristics

Maturity: early / 7,5

Light sprout colour: red violet / RV

Flower colour: light purple / LP

Number of berries: very high / 4

Foliage - initial development: quite fast / 7

Foliage - final development: average / 7

Foliage - firmness: quite fragile / 5,5

Skin colour: red / R

Flesh colour: yellow / 7

Skin finish: excellent / 8

Tuber shape: long / L

Regularity of tuber shape: regular / 7

Tuberisation: high / 9

Tuber size: average / 6

Regularity of size: quite regular / 6

Dormancy: average / 6

Sensitivity and quality

Secondary growth: quite sensitive / 6,5

Black spot (internal bruising): slightly sensitive / 7

Growth cracks: sensitive / 5

Harvest damage: slightly sensitive / 7

Metribuzin - pre-emergence: sensitive / 5

Metribuzin - post-emergence: sensitive / 5

Bentazon: slightly sensitive / 7

Ethylene: - / -

Underwater weight: 391

Dry matter content: 21,3%

Cooking type: firm / A

Discolouration (after cooking): slightly sensitive / 7

Raw discolouration: sensitive / 5,5

Fries quality: unsuitable / 5

Dry matter distribution: excellent / 8

Crisping quality: unsuitable / 4,5

TGA-level: 5,5



How to grow

www.agricopotatoes.com

Fertilization ¹

Nitrogen: standard advice - 10%.

Phosphate: standard advice.

Potassium: standard advice.

Seed treatment

Can be planted directly out of the cold store, preferably after a short heat shock. Desprouting more than once can have a negative impact on yield and size regularity.

Planting distance

30/50 mm: 20 cm (67.000 plants/ha)

Weeds

Do not use metribuzin (Sencor), as it gives a severe foliage reaction and a serious yield reduction.

Fungicides

A strict preventive spraying schedule against late blight is necessary.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

Storage - .

Preferred storage temperature is 4 °C. Suitable for medium to long term storage.

¹ Level of fertilization is based on soil analysis.



Say potato,
say **Agrico.**

Resistances

Virus - Leaf roll: - / -

Virus - Yn: slightly to not susceptible / 8

Virus - Yntn (tuber): slightly susceptible / 7

Spraing: susceptible / 5

Late blight - foliage: very susceptible / 4

Late blight - tuber: slightly susceptible / 7

Fusarium: slightly susceptible / 6

Common scab: slightly susceptible / 6

Powdery scab: slightly susceptible / 6

Black dot: slightly susceptible / 7

Wart disease F1: - / -

Wart disease F6: - / -

Wart disease F18: - / -

Golden nematode Ro 1/4: - / -

Golden nematode Ro 2/3: - / -

White nematode Pa 2: - / -

White nematode Pa 3: - / -