

# Ditta.

A versatile, early maincrop ware and salad potato, suitable for conventional and organic cultivation.




Organic



Retail  
Fresh  
Salads

- ▶ **Excellent consumption quality**
- ▶ **Good storability**
- ▶ **Multiple resistances**



## characteristics

### General characteristics

Breeder: Nieder Österreichische  
Saatbaugenossenschaft

Crossing: BINTJE x QUARTA

Breedersright: Expired

Seedling: SW 80-0540

### Plant and tuber characteristics

Maturity: early maincrop / 6

Light sprout colour: blue violet / BV

Flower colour: white / W

Number of berries: very high / 4

Foliage - initial development: quite fast / 7,5

Foliage - final development: good / 8

Foliage - firmness: quite firm / 7

Skin colour: yellow / Y

Flesh colour: light yellow / 6,5

Skin finish: good / 7

Tuber shape: long oval / LO

Regularity of tuber shape: regular / 7

Tuberisation: quite high / 7

Tuber size: quite large / 7

Regularity of size: quite regular / 6,5

Dormancy: quite long / 7

### Sensitivity and quality

Secondary growth: quite sensitive / 6

Black spot (internal bruising): slightly sensitive / 7

Growth cracks: quite sensitive / 6,5

Harvest damage: sensitive / 5,5

Metribuzin - pre-emergence: slightly sensitive / 7

Metribuzin - post-emergence: slightly sensitive / 7

Ethylene: - / -

Underwater weight: 369

Dry matter content: 20,2%

Cooking type: firm / A

Discolouration (after cooking): quite sensitive / 6

Raw discolouration: very sensitive / 4

Fries quality: below average / 5,5

Dry matter distribution: - / -

Crisping quality: unsuitable / 4

TGA-level: -

# How to grow



[www.agricopotatoes.com](http://www.agricopotatoes.com)

## Fertilization <sup>1</sup>

**Nitrogen:** standard advice - 10%.

**Phosphate:** standard advice.

**Potassium:** standard advice.

## Seed treatment

Can be planted directly out of the cold store, preferably after a short heat shock. Desprouting more than once can have a negative impact on yield and size regularity.

## Planting distance

28/35 mm: 22 cm (60.500 plants/ha)

35/50 mm: 27 cm (50.000 plants/ha)

## Weeds

A standard use of metribuzin (Sencor), before or after emerge, can give some foliage reaction, but rarely any yield reduction.

## Fungicides

A strict preventive spraying schedule against late blight is necessary.

## Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

## Storage - .

Preferred storage temperature is 5 °C. Suitable for medium term storage.

<sup>1</sup> Level of fertilization is based on soil analysis.



Say potato,  
say **Agrico.**

## Resistances

**Virus - Leaf roll:** slightly susceptible / 7

**Virus - Yn:** slightly susceptible / 7,5

**Virus - Yntn (tuber):** susceptible / 5

**Spraing:** susceptible / 5

**Late blight - foliage:** susceptible / 5,5

**Late blight - tuber:** slightly susceptible / 7

**Fusarium:** slightly susceptible / 6

**Common scab:** slightly susceptible / 6

**Powdery scab:** slightly susceptible / 6,5

**Black dot:** slightly susceptible / 6

**Wart disease F1:** resistant / 10

**Wart disease F6:** - / -

**Wart disease F18:** - / -

**Golden nematode Ro 1/4:** resistant / 9

**Golden nematode Ro 2/3:** - / -

**White nematode Pa 2:** - / -

**White nematode Pa 3:** - / -