

Levante.

An early maincrop, high yielding ware potato variety with a nice yellow skin and resilient to late blight in foliage and tuber.



Organic



Retail
Fresh



Traditional

- ▶ **Late blight resistance**
- ▶ **Beautiful skin**
- ▶ **Strong plant**
- ▶ **Multiple resistances**
- ▶ **High tuberization**
- ▶ **High yield**



characteristics

General characteristics

Breeder: Agrico Research B.V.
 Crossing: AR 01-3218 x ALMERA
 Breedersright: 31-12-2048
 Seedling: AR 07-3009

Plant and tuber characteristics

Maturity: early maincrop / 5,5
 Light sprout colour: green red / GR
 Flower colour: white / W
 Number of berries: above average / 5,5
 Foliage - initial development: quite fast / 7
 Foliage - final development: good / 8
 Foliage - firmness: quite firm / 7
 Skin colour: yellow / Y
 Flesh colour: light yellow / 6
 Skin finish: good / 7
 Tuber shape: long oval / LO
 Regularity of tuber shape: quite regular / 6,5
 Tuberisation: high / 8
 Tuber size: quite large / 7
 Regularity of size: regular / 7
 Dormancy: average / 6,5

Sensitivity and quality

Secondary growth: slightly sensitive / 7
 Black spot (internal bruising): quite sensitive / 6,5
 Growth cracks: slightly sensitive / 7
 Harvest damage: quite sensitive / 6
 Metribuzin - pre-emergence: quite sensitive / 6,5
 Metribuzin - post-emergence: sensitive / 5,5
 Ethylene: 5% fewer tubers / 4
 Underwater weight: 365
 Dry matter content: 19,9%
 Cooking type: rather floury / B
 Discolouration (after cooking): slightly sensitive / 7
 Raw discolouration: sensitive / 5,5
 Fries quality: below average / 5,5
 Dry matter distribution: - / -
 Crisping quality: unsuitable / 4,5
 TGA-level: 4,73

How to grow



www.agricopotatoes.com

Fertilization ¹

Nitrogen: standard advice. Preferably as split application, second treatment shortly after tuberization.

Phosphate: standard advice.

Potassium: standard advice.

Seed treatment

Preferably plant the tubers into warm soil, as the initial development of this variety is slow. Desprouting can have a negative impact on yield and size regularity. A short heat shock or proper pre-sprouting is recommended.

Planting distance

28/35 mm: 22 cm (60.500 plants/ha)

35/55 mm: 27 cm (49.500 plants/ha)

Weeds

A standard use of metribuzin (Sencor), before or after emerge, doesn't give foliage reaction or yield reduction.

Fungicides

Despite the late blight resistance, remain alert for infections.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least three weeks after haulm killing before harvest.

Storage - .

Preferred storage temperature is 4 °C. Suitable for medium to long term storage.

¹ Level of fertilization is based on soil analysis.



Say potato,
say **Agrico.**

Resistances

Virus - Leaf roll: - / -

Virus - Yn: slightly susceptible / 6

Virus - Yntn (tuber): slightly to not susceptible / 8,5

Spraing: very susceptible / 4,5

Late blight - foliage: slightly to not susceptible / 8,5

Late blight - tuber: slightly susceptible / 7,5

Fusarium: slightly susceptible / 6

Common scab: susceptible / 5

Powdery scab: slightly susceptible / 6,5

Black dot: slightly susceptible / 7,5

Wart disease F1: - / -

Wart disease F6: - / -

Wart disease F18: - / -

Golden nematode Ro 1/4: resistant / 9

Golden nematode Ro 2/3: - / -

White nematode Pa 2: - / -

White nematode Pa 3: - / -