


Napoleon.



Processing
Crisps

Maincrop crisping potato variety with a good uniformity of size and a high dry matter content.

- ▶ **Good crisping quality**
- ▶ **Excellent storability**
- ▶ **Good virus resistances**



characteristics

General characteristics

Breeder: Agrico Research B.V.
 Crossing: ANTINA x ALICANTE
 Breedersright: 31-12-2051
 Seedling: AR 10-4637

Plant and tuber characteristics

Maturity: maincrop / 5
 Light sprout colour: red violet / RV
 Flower colour: white / W
 Number of berries: very high / 4
 Foliage - initial development: quite fast / 7
 Foliage - final development: good / 8
 Foliage - firmness: quite firm / 7,5
 Skin colour: yellow / Y
 Flesh colour: light yellow / 6,5
 Skin finish: quite poor / 5
 Tuber shape: round oval / RO
 Regularity of tuber shape: regular / 7
 Tuberisation: average / 6
 Tuber size: quite large / 7,5
 Regularity of size: regular / 7
 Dormancy: average / 6

Sensitivity and quality

Secondary growth: quite sensitive / 6,5
 Black spot (internal bruising): sensitive / 5,5
 Growth cracks: quite sensitive / 6
 Harvest damage: very sensitive / 4,5
 Metribuzin - pre-emergence: slightly sensitive / 7
 Metribuzin - post-emergence: slightly sensitive / 7
 Ethylene: 5% fewer to 5% more tubers / 5
 Underwater weight: 450
 Dry matter content: 24,1%
 Cooking type: floury / BC
 Discolouration (after cooking): quite sensitive / 6
 Raw discolouration: quite sensitive / 6
 Fries quality: good / 7
 Dry matter distribution: good / 7
 Crisping quality: excellent / 8
 TGA-level: 4,0

How to grow



www.agricopotatoes.com

Fertilization ¹

Nitrogen: standard advice - 10%.

Phosphate: standard advice.

Potassium: standard advice + 20%.

Seed treatment

Proper pre-sprouting after taking off the top sprout can give a positive result on the tuber set. Desprouting can have a negative impact on yield and size regularity.

Planting distance

28/35 mm: 25 cm (54.000 plants/ha)

35/55 mm: 30 cm (45.000 plants/ha)

50/55 mm: 40 cm (33.000 plants/ha)

Weeds

A standard use of metribuzin (Sencor), before or after emerge, doesn't give foliage reaction or yield reduction.

Fungicides

A preventive spraying schedule against late blight is advised.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least three weeks after haulm killing before harvest.

Storage - .

For processing purposes, do not store below 9 °C. Suitable for medium to long term storage.

¹ Level of fertilization is based on soil analysis.



Say potato,
say Agrico.

Resistances

Virus - Leaf roll: - / -

Virus - Yn: slightly to not susceptible / 8

Virus - Yntn (tuber): slightly to not susceptible / 8

Spraing: susceptible / 5

Late blight - foliage: slightly susceptible / 6

Late blight - tuber: slightly susceptible / 6,5

Fusarium: slightly susceptible / 6,5

Common scab: slightly susceptible / 6

Powdery scab: - / -

Black dot: slightly susceptible / 6

Wart disease F1: resistant / 10

Wart disease F6: - / -

Wart disease F18: - / -

Golden nematode Ro 1/4: resistant / 9

Golden nematode Ro 2/3: - / -

White nematode Pa 2: susceptible / 4

White nematode Pa 3: susceptible / 2