

Vogue.



Traditional

Second early ware potato variety with uniformly sized, big tubers.

- ▶ **Rather firm cooking**
- ▶ **Large tubers**
- ▶ **Early tuberisation**



characteristics

General characteristics

Breeder: Konst Research B.V.
 Crossing: MARABEL x AMOROSA
 Breedersright: 31-12-2046
 Seedling: KO 03-2027

Plant and tuber characteristics

Maturity: second early / 6,5
 Light sprout colour: green red / GR
 Flower colour: white / W
 Number of berries: very high / 4
 Foliage - initial development: quite fast / 7
 Foliage - final development: quite good / 7,5
 Foliage - firmness: quite firm / 7
 Skin colour: yellow / Y
 Flesh colour: yellow / 7
 Skin finish: average / 6
 Tuber shape: oval long / OL
 Regularity of tuber shape: regular / 7
 Tuberisation: quite low / 5,5
 Tuber size: large / 8
 Regularity of size: regular / 7
 Dormancy: quite short / 5

Sensitivity and quality

Secondary growth: quite sensitive / 6,5
 Black spot (internal bruising): slightly to not sensitive / 8
 Growth cracks: quite sensitive / 6,5
 Harvest damage: quite sensitive / 6
 Metribuzin - pre-emergence: slightly to not sensitive / 8
 Metribuzin - post-emergence: slightly sensitive / 7,5
 Ethylene: 5% fewer to 5% more tubers / 5
 Underwater weight: 311
 Dry matter content: 17,3%
 Cooking type: rather firm / AB
 Discolouration (after cooking): slightly sensitive / 7,5
 Raw discolouration: - / -
 Fries quality: unsuitable / 5
 Dry matter distribution: - / -
 Crisping quality: unsuitable / 4
 TGA-level: 3,7

How to grow



www.agricopotatoes.com

Fertilization ¹

Nitrogen: standard advice - 10%. Preferably as split application, second treatment shortly after tuberization.

Phosphate: standard advice.

Potassium: standard advice + 10%.

Seed treatment

Proper pre-sprouting after taking off the top sprout can give a positive result on the tuber set. Desprouting more than once can have a negative impact on yield and size regularity.

Planting distance

28/35 mm: 22 cm (60.500 plants/ha)

35/55 mm: 27 cm (50.000 plants/ha)

Weeds

A standard use of metribuzin (Sencor), before or after emerge, doesn't give foliage reaction or yield reduction.

Fungicides

A strict preventive spraying schedule against late blight is necessary.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least three weeks after haulm killing before harvest.

Storage - .

Preferred storage temperature is 5 °C. Suitable for delivery straight from the field or short term storage.

¹ Level of fertilization is based on soil analysis.



Say potato,
say Agrico.

Resistances

Virus - Leaf roll: - / -

Virus - Yn: slightly susceptible / 6,5

Virus - Yntn (tuber): susceptible / 5

Spraing: very susceptible / 4

Late blight - foliage: very susceptible / 4

Late blight - tuber: slightly susceptible / 6

Fusarium: slightly susceptible / 6,5

Common scab: slightly susceptible / 6

Powdery scab: very susceptible / 2

Black dot: very susceptible / 4,5

Wart disease F1: slightly susceptible / 9

Wart disease F6: - / -

Wart disease F18: - / -

Golden nematode Ro 1/4: resistant / 9

Golden nematode Ro 2/3: - / -

White nematode Pa 2: - / -

White nematode Pa 3: - / -