

# Vogue.

Second early ware potato variety with uniformly sized, big tubers.



Traditional

- ▶ **Rather firm cooking**
- ▶ **Large tubers**
- ▶ **Early tuberisation**



## characteristics

### General characteristics

Breeder: Konst Research B.V.  
 Crossing: MARABEL x AMOROSA  
 Breedersright: 31-12-2046  
 Seedling: KO 03-2027

### Plant and tuber characteristics

Maturity: second early / 6,5  
 Light sprout colour: green red / GR  
 Flower colour: white / W  
 Number of berries: very high / 4  
 Foliage - initial development: quite fast / 7  
 Foliage - final development: quite good / 7,5  
 Foliage - firmness: quite firm / 7  
 Skin colour: yellow / Y  
 Flesh colour: yellow / 7  
 Skin finish: average / 6  
 Tuber shape: oval long / OL  
 Regularity of tuber shape: regular / 7  
 Tuberisation: quite low / 5,5  
 Tuber size: large / 8  
 Regularity of size: regular / 7  
 Dormancy: quite short / 5

### Sensitivity and quality

Secondary growth: quite sensitive / 6,5  
 Black spot (internal bruising): slightly to not sensitive / 8  
 Growth cracks: quite sensitive / 6,5  
 Harvest damage: quite sensitive / 6  
 Metribuzin - pre-emergence: slightly to not sensitive / 8  
 Metribuzin - post-emergence: slightly sensitive / 7,5  
 Bentazon: slightly sensitive / 7  
 Ethylene: 5% fewer to 5% more tubers / 5  
 Underwater weight: 311  
 Dry matter content: 17,3%  
 Cooking type: rather firm / AB  
 Discolouration (after cooking): slightly sensitive / 7,5  
 Raw discolouration: - / -  
 Fries quality: unsuitable / 5  
 Dry matter distribution: - / -  
 Crisping quality: unsuitable / 4  
 TGA-level: 3,7

# How to grow

## Fertilization <sup>1</sup>

**Nitrogen:** standard advice - 10%. Preferably as split application, second treatment shortly after tuberization.

**Phosphate:** standard advice.

**Potassium:** standard advice + 10%.

## Seed treatment

Proper pre-sprouting after taking off the top sprout can give a positive result on the tuber set. Desprouting more than once can have a negative impact on yield and size regularity.

## Planting distance

28/35 mm: 22 cm (60.500 plants/ha)

35/55 mm: 27 cm (50.000 plants/ha)



[www.agricopotatoes.com](http://www.agricopotatoes.com)

## Weeds

A standard use of metribuzin (Sencor), before or after emerge, doesn't give foliage reaction or yield reduction.

## Fungicides

A strict preventive spraying schedule against late blight is necessary.

## Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least three weeks after haulm killing before harvest.

## Storage - .

Preferred storage temperature is 5 °C. Suitable for delivery straight from the field or short term storage.

<sup>1</sup> Level of fertilization is based on soil analysis.



**Say potato,  
say Agrico.**

## Resistances

**Virus - Leaf roll:** - / -

**Virus - Yn:** slightly susceptible / 6,5

**Virus - Yntn (tuber):** susceptible / 5

**Spraying:** very susceptible / 4

**Late blight - foliage:** very susceptible / 4

**Late blight - tuber:** slightly susceptible / 6

**Fusarium:** slightly susceptible / 6,5

**Common scab:** slightly susceptible / 6

**Powdery scab:** very susceptible / 2

**Black dot:** very susceptible / 4,5

**Wart disease F1:** slightly susceptible / 9

**Wart disease F6:** - / -

**Wart disease F18:** - / -

**Golden nematode Ro 1/4:** resistant / 9

**Golden nematode Ro 2/3:** - / -

**White nematode Pa 2:** - / -

**White nematode Pa 3:** - / -