CULTIVATION RECOMMENDATION

Vogue.

Second early ware potato variety with uniformly sized, big tubers.

Traditional

Image:

- **Rather firm cooking**
- **Large tubers**
- **Early tuberisation**

characteristics

General characteristics

Breeder: Konst Research B.V.
Crossing: MARABEL x AMOROSA
Breedersright: 31-12-2046
Seedling: KO 03-2027

Plant and tuber characteristics

Maturity: second early / 6,5
Light sprout colour: green red / GR
Flower colour: white / W
Number of berries: very high / 4
Foliage - initial development: quite fast / 7
Foliage - final development: quite good / 7,5
Foliage - firmness: quite firm / 7
Skin colour: yellow / Y
Flesh colour: yellow / 7
Skin finish: average / 6
Tuber shape: oval long / OL
Regularity of tuber shape: regular / 7
Tuberisation: quite low / 5,5
Tuber size: large / 8
Regularity of size: regular / 7
Dormancy: quite short / 5

Sensitivity and quality

Secondary growth: quite sensitive / 6,5
Black spot (internal bruising): slightly to not sensitive / 8
Growth cracks: quite sensitive / 6,5
Harvest damage: quite sensitive / 6
Metribuzin - pre-emergence: slightly to not sensitive / 8
Metribuzin - post-emergence: slightly sensitive / 7,5
Ethylene: 5% fewer to 5% more tubers / 5
Underwater weight: 311
Dry matter content: 17,3%
Cooking type: rather firm / AB
Discolouration (after cooking): slightly sensitive / 7,5
Raw discolouration: - / -
Fries quality: unsuitable / 5
Dry matter distribution: - / -
Crisping quality: unsuitable / 4
TGA-level: 3,7 mg/100g

Agrico has taken all reasonable care in producing this variety description. However, please note that any information may be subject to change and Agrico accepts no responsibility for errors or omissions. In no event shall Agrico be liable for damages arising out of or related to the information contained within this description.

AGRICO

CULTIVATION RECOMMENDATION

How to grow

www.agricopotatoes.com

Fertilization [1]

Nitrogen: standard advice - 10%.Preferably as split application, second treatment shortly after tuberization.

Phosphate: standard advice.

Potassium: standard advice + 10%.

Seed treatment

Proper pre-sprouting after taking off the top sprout can give a positive result on the tuber set.Desprouting more than once can have a negative impact on yield and size regularity.

Planting distance

28/35 mm: 25 cm (53.000 plants/ha)
35/55 mm: 30 cm (44.500 plants/ha)

Weeds

A standard use of metribuzin (Sencor), before or after emerge, doesn't give foliage reaction or yield reduction.

Fungicides

A strict preventive spraying schedule against late blight is necessary.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least three weeks after haulm killing before harvest.

Storage - .

Preferred storage temperature is 5 ºC. Suitable for delivery straight from the field or short term storage.

[1] Level of fertilization is based on soil analysis.



Say potato, say Agrico.

Resistances

Virus - Leaf roll: - / -
Virus - Yn: slightly susceptible / 6,5
Virus - Yntn (tuber): susceptible / 5
Spraing: very susceptible / 4
Late blight - foliage: very susceptible / 4
Late blight - tuber: slightly susceptible / 6
Fusarium: slightly susceptible / 6,5
Common scab: slightly susceptible / 6

Powdery scab: very susceptible / 2
Black dot: very susceptible / 4,5
Wart disease F1: slightly susceptible / 9
Wart disease F6: - / -
Wart disease F18: - / -
Golden nematode Ro 1/4: resistant / 9
Golden nematode Ro 2/3: - / -
White nematode Pa 2: - / -
White nematode Pa 3: - / -

Agrico • Duit 15 • 8305 BB Emmeloord • The Netherlands +31 (0)527 63 99 11 info@agrico.nl www.agricopotatoes.com