
CULTIVATION RECOMMENDATION

Fajah.

A strong ware potato variety with regular large tubers with a relatively high dry matter content.

- **Large tubers and high yield**
- **Long oval tubers and uniform in size**
- **Short dormancy, suitable for second crop**

Traditional



characteristics

General characteristics

Breeder: R. Rispens
Crossing: LANORMA x AMBITION
Breedersright: 31-12-2052
Seedling: RP 10-1003

Plant and tuber characteristics

Maturity: early maincrop / 5,5
Light sprout colour: red / R
Flower colour: white / W
Number of berries: below average / 7
Foliage - initial development: fast / 8
Foliage - final development: good / 8
Foliage - firmness: quite firm / 7,5
Skin colour: yellow / Y
Flesh colour: light yellow / 6
Skin finish: above average / 6,5
Tuber shape: long oval / LO
Regularity of tuber shape: regular / 7
Tuberisation: average / 6
Tuber size: large / 8
Regularity of size: quite regular / 6,5
Dormancy: average / 6

Sensitivity and quality

Secondary growth: slightly sensitive / 7
Black spot (internal bruising): slightly sensitive / 7
Growth cracks: quite sensitive / 6,5
Harvest damage: sensitive / 5
Metribuzin - pre-emergence: slightly sensitive / 7
Metribuzin - post-emergence: quite sensitive / 6,5
Ethylene: 15% to 25% more tubers / 7
Underwater weight: 354
Dry matter content: 19,4%
Cooking type: floury / BC
Discolouration (after cooking): quite sensitive / 6,5
Raw discolouration: quite sensitive / 6
Fries quality: good / 7
Dry matter distribution: - / -
Crisping quality: above average / 6,5
TGA-level: 4,5 mg/100g

Agrico has taken all reasonable care in producing this variety description. However, please note that any information may be subject to change and Agrico accepts no responsibility for errors or omissions. In no event shall Agrico be liable for damages arising out of or related to the information contained within this description.

AGRICO

CULTIVATION RECOMMENDATION

How to grow

www.agricopotatoes.com

Fertilization [1]

Nitrogen: standard advice + 10%.
Phosphate: standard advice.
Potassium: standard advice.

Seed treatment

Desprouting can have a negative impact on yield and size regularity.

Planting distance

28/35 mm: 27 cm (50.000 plants/ha)
35/55 mm: 33 cm (40.000 plants/ha)

Weeds

A standard use of metribuzin (Sencor), before or after emerge, can give some foliage reaction and some yield reduction.

Fungicides

A strict preventive spraying schedule against late blight is necessary.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

Storage - .

Preferred storage temperature is 5 ºC. Suitable for medium term storage.

[1] Level of fertilization is based on soil analysis.



Say potato, say Agrico.

Resistances

Virus - Leaf roll: - / -
Virus - Yn: susceptible / 5,5
Virus - Yntn (tuber): slightly susceptible / 6,5
Spraing: susceptible / 5
Late blight - foliage: very susceptible / 4
Late blight - tuber: susceptible / 5,5
Fusarium: susceptible / 5,5
Common scab: susceptible / 5

Powdery scab: - / -
Black dot: susceptible / 5,5
Wart disease F1: very susceptible* / 1*
Wart disease F6: - / -
Wart disease F18: - / -
Golden nematode Ro 1/4: resistant / 9
Golden nematode Ro 2/3: - / -
White nematode Pa 2: susceptible / 3
White nematode Pa 3: susceptible / 2

Agrico • Duit 15 • 8305 BB Emmeloord • The Netherlands +31 (0)527 63 99 11 info@agrico.nl www.agricopotatoes.com