


# Fajah.



Traditional

A strong ware potato variety with regular large tubers with a relatively high dry matter content.

- ▶ **Large tubers and high yield**
- ▶ **Long oval tubers and uniform in size**
- ▶ **Short dormancy, suitable for second crop**



## characteristics

### General characteristics

Breeder: R. Rispens

Crossing: LANORMA x AMBITION

Breedersright: 31-12-2052

Seedling: RP 10-1003

### Plant and tuber characteristics

Maturity: early maincrop / 5,5

Light sprout colour: red / R

Flower colour: white / W

Number of berries: below average / 7

Foliage - initial development: fast / 8

Foliage - final development: good / 8

Foliage - firmness: quite firm / 7,5

Skin colour: yellow / Y

Flesh colour: light yellow / 6

Skin finish: above average / 6,5

Tuber shape: long oval / LO

Regularity of tuber shape: regular / 7

Tuberisation: average / 6

Tuber size: quite large / 7,5

Regularity of size: quite regular / 6,5

Dormancy: quite short / 5,5

### Sensitivity and quality

Secondary growth: quite sensitive / 6,5

Black spot (internal bruising): slightly sensitive / 7

Growth cracks: quite sensitive / 6,5

Harvest damage: sensitive / 5

Metribuzin - pre-emergence: slightly sensitive / 7

Metribuzin - post-emergence: quite sensitive / 6,5

Ethylene: 15% to 25% more tubers / 7

Underwater weight: 354

Dry matter content: 19,4%

Cooking type: floury / BC

Discolouration (after cooking): quite sensitive / 6,5

Raw discolouration: quite sensitive / 6

Fries quality: good / 7

Dry matter distribution: - / -

Crisping quality: above average / 6,5

TGA-level: 4,5

# How to grow



[www.agricopotatoes.com](http://www.agricopotatoes.com)

## Fertilization <sup>1</sup>

**Nitrogen:** standard advice + 10%.

**Phosphate:** standard advice.

**Potassium:** standard advice.

## Seed treatment

Desprouting can have a negative impact on yield and size regularity.

## Planting distance

28/35 mm: 27 cm (50.000 plants/ha)

35/55 mm: 33 cm (40.000 plants/ha)

## Weeds

A standard use of metribuzin (Sencor), before or after emerge, can give some foliage reaction and some yield reduction.

## Fungicides

A strict preventive spraying schedule against late blight is necessary.

## Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

## Storage - .

Preferred storage temperature is 5 °C. Suitable for medium term storage.

<sup>1</sup> Level of fertilization is based on soil analysis.



Say potato,  
say Agrico.

## Resistances

**Virus - Leaf roll:** - / -

**Virus - Yn:** susceptible / 5,5

**Virus - Yntn (tuber):** slightly susceptible / 6,5

**Spraing:** susceptible / 5

**Late blight - foliage:** very susceptible / 4

**Late blight - tuber:** susceptible / 5,5

**Fusarium:** susceptible / 5,5

**Common scab:** susceptible / 5

**Powdery scab:** - / -

**Black dot:** susceptible / 5,5

**Wart disease F1:** very susceptible\* / 1\*

**Wart disease F6:** - / -

**Wart disease F18:** - / -

**Golden nematode Ro 1/4:** resistant / 9

**Golden nematode Ro 2/3:** - / -

**White nematode Pa 2:** susceptible / 3

**White nematode Pa 3:** susceptible / 2