

Riviera.



Traditional

A first early ware potato variety with an attractive skin, excellent heat tolerance, good virus resistance and early a good yield.

- ▶ **Large tubers**
- ▶ **Beautiful appearance**
- ▶ **Good virus resistances**



characteristics

General characteristics

Breeder: Mts. Boerhave

Crossing: MINERVA x ALCMARIA

Breedersright: 31-12-2030

Seedling: BH 86-0266

Plant and tuber characteristics

Maturity: first early / 8

Light sprout colour: blue violet / BV

Flower colour: white / W

Number of berries: very high / 4

Foliage - initial development: quite fast / 7

Foliage - final development: average / 7

Foliage - firmness: average / 6,5

Skin colour: yellow / Y

Flesh colour: light yellow / 6

Skin finish: excellent / 8

Tuber shape: oval round / OR

Regularity of tuber shape: quite regular / 6

Tuberisation: quite low / 5

Tuber size: quite large / 7,5

Regularity of size: regular / 7

Dormancy: average / 6

Sensitivity and quality

Secondary growth: quite sensitive / 6,5

Black spot (internal bruising): slightly to not sensitive / 8

Growth cracks: quite sensitive / 6,5

Harvest damage: quite sensitive / 6

Metribuzin - pre-emergence: slightly to not sensitive / 8

Metribuzin - post-emergence: slightly to not sensitive / 8

Ethylene: 5% to 15% more tubers / 6

Underwater weight: 318

Dry matter content: 17,7%

Cooking type: rather firm / AB

Discolouration (after cooking): sensitive / 5,5

Raw discolouration: - / -

Fries quality: unsuitable / 4,5

Dry matter distribution: - / -

Crisping quality: unsuitable / 4

TGA-level: 4,8

How to grow



www.agricopotatoes.com

Fertilization ¹

Nitrogen: standard advice + 10%.

Phosphate: standard advice.

Potassium: standard advice.

Seed treatment

Proper pre-sprouting after taking off the top sprout can give a positive result on the tuber set. Desprouting more than once can have a negative impact on yield and size regularity.

Planting distance

28/35 mm: 22 cm (60.500 plants/ha)

35/55 mm: 27 cm (50.000 plants/ha)

Weeds

A standard use of metribuzin (Sencor), before or after emerge, doesn't give foliage reaction or yield reduction.

Fungicides

A strict preventive spraying schedule against late blight is necessary.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

Storage - .

Preferred storage temperature is 4 °C. Suitable for delivery straight from the field or short term storage.

¹ Level of fertilization is based on soil analysis.



Say potato,
say **Agrico.**

Resistances

Virus - Leaf roll: slightly susceptible / 7

Virus - Yn: slightly to not susceptible / 8

Virus - Yntn (tuber): slightly susceptible / 7

Spraing: very susceptible / 4

Late blight - foliage: very susceptible / 3

Late blight - tuber: susceptible / 5

Fusarium: slightly susceptible / 6

Common scab: susceptible / 5,5

Powdery scab: very susceptible / 2

Black dot: susceptible / 5,5

Wart disease F1: resistant / 10

Wart disease F6: - / -

Wart disease F18: - / -

Golden nematode Ro 1/4: resistant / 9

Golden nematode Ro 2/3: - / -

White nematode Pa 2: - / -

White nematode Pa 3: - / -