

Sante.

Early maincrop ware potato with a good yield.

- ▶ **High dry matter content**
- ▶ **Rather floury**
- ▶ **Multiple resistances**



**Retail
Fresh**



Traditional



characteristics

General characteristics

Breeder: J. Vegter

Crossing: WY 66-13-636 x AM 66-0042

Breedersright: Expired

Seedling: VE 72-13-225

Plant and tuber characteristics

Maturity: early maincrop / 6

Light sprout colour: red violet / RV

Flower colour: white / W

Number of berries: very high / 4

Foliage - initial development: quite fast / 7

Foliage - final development: quite good / 7,5

Foliage - firmness: quite firm / 7

Skin colour: yellow / Y

Flesh colour: light yellow / 6

Skin finish: average / 6

Tuber shape: oval round / OR

Regularity of tuber shape: quite regular / 6

Tuberisation: quite high / 7

Tuber size: quite large / 7

Regularity of size: quite regular / 6,5

Dormancy: average / 6

Sensitivity and quality

Secondary growth: quite sensitive / 6,5

Black spot (internal bruising): slightly sensitive / 7,5

Growth cracks: quite sensitive / 6,5

Harvest damage: quite sensitive / 6

Metribuzin - pre-emergence: quite sensitive / 6

Metribuzin - post-emergence: slightly sensitive / 7

Ethylene: - / -

Underwater weight: 425

Dry matter content: 22,9%

Cooking type: rather floury / B

Discolouration (after cooking): sensitive / 5,5

Raw discolouration: - / -

Fries quality: average / 6

Dry matter distribution: - / -

Crisping quality: unsuitable / 5

TGA-level: 3,9

How to grow



www.agricopotatoes.com

Fertilization ¹

Nitrogen: standard advice.

Phosphate: standard advice.

Potassium: standard advice.

Seed treatment

Proper pre-sprouting after taking off the top sprout can give a positive result on the tuber set. Desprouting more than once can have a negative impact on yield and size regularity.

Planting distance

28/35 mm: 25 cm (54.000 plants/ha)

35/55 mm: 30 cm (45.000 plants/ha)

Weeds

A standard use of metribuzin (Sencor), before or after emerge, can give some foliage reaction and some yield reduction.

Fungicides

A strict preventive spraying schedule against late blight is necessary.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

Storage - .

Preferred storage temperature is 5 °C. Suitable for medium term storage.

¹ Level of fertilization is based on soil analysis.



Say potato,
say **Agrico.**

Resistances

Virus - Leaf roll: slightly susceptible / 7

Virus - Yn: slightly to not susceptible / 8,5

Virus - Yntn (tuber): slightly to not susceptible / 9

Spraing: slightly susceptible / 6

Late blight - foliage: susceptible / 5

Late blight - tuber: slightly susceptible / 7

Fusarium: slightly susceptible / 6

Common scab: susceptible / 5

Powdery scab: slightly to not susceptible / 8

Black dot: susceptible / 5,5

Wart disease F1: resistant / 10

Wart disease F6: - / -

Wart disease F18: - / -

Golden nematode Ro 1/4: resistant / 9

Golden nematode Ro 2/3: field resistant / 7

White nematode Pa 2: field resistant / 7

White nematode Pa 3: susceptible / 3